


Friday, March 6, 2020

## 5.5 The Derivative of $y = \tan x$



### Bellwork:

Use the quotient identity for  $\tan x$  to find the derivative of  $y = \tan x$ .

$$y = \frac{\sin x}{\cos x}$$

Use the quotient rule.

$$\begin{aligned} y' &= \frac{(\cos x)(\cos x) - (-\sin x)\sin x}{\cos^2 x} \\ &= \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\ &= \frac{1}{\cos^2 x} \\ &= \sec^2 x \end{aligned}$$

Do you ever HAVE to use the derivative of  $\tan x$ ? Why or why not?

No,  $\tan x$  can always be expressed in terms of  $\sin x$  and  $\cos x$ .

When will the derivative of  $\tan x$  be undefined? What type of discontinuity occurs here?

When  $\cos^2 x = 0$ .

Vertical asymptote.



Differentiate each of the following functions:

$$1) f(x) = \tan^2(2x)$$

$$f(x) = (\tan 2x)^2$$

$$f'(x) = 2(\tan 2x)(\sec^2 2x)(2)$$

$$= 4 \tan 2x \sec^2 2x$$

$$2) g(x) = -3x^2 + \sin x - 4 \tan 2x$$

$$g'(x) = 6x^{-2} + \cos x - 4(\sec^2 2x)(2)$$

$$= \frac{6}{x^3} + \cos x - 8 \sec^2 2x$$

$$3) h(x) = \underbrace{(x-5)^2}_{1^{st}} \underbrace{(\cos^2 x + 2 \tan x)}_{2^{nd}}$$

$$h'(x) = 2(x-5)(\cos^2 x + 2 \tan x) + (x-5)^2(-2 \cos x \sin x + 2 \sec^2 x)$$

$$= 2(x-5)(\cos^2 x + 2 \tan x) - 2(x-5)(\cos x \sin x - \sec^2 x)$$

$$= 2(x-5)[(\cos^2 x + 2 \tan x) - (x-5)(\cos x \sin x - \sec^2 x)]$$

$$4) y = \sin 2x \tan 2x \cos^2 2x$$

$$y = (\sin 2x) \left( \frac{\sin 2x}{\cos 2x} \right) (\cos^2 2x)$$

$$y = \underbrace{\sin^2 2x}_{1^{st}} \cos 2x$$

$$\sin 2x = 2 \sin x \cos x$$

$$\frac{1}{2} \sin 2x = \sin x \cos x$$

Remember that you can use trig identities to write expressions in simpler forms sometimes! Please do that, even if the text book does not. For example, how can you simplify  $f(x) = \tan x \cos x$ ?  $g(x) = \cot x \sec x$ ?

$$f(x) = \tan x \cos x \quad g(x) = \cot x \sec x$$

$$f(x) = \frac{\sin x}{\cos x} \cos x = \frac{\cos x}{\sin x} \cdot \frac{1}{\cos x}$$

$$f(x) = \sin x \quad = \frac{1}{\sin x}$$

$$f'(x) = \cos x \quad g'(x) = \frac{0 - \cos x}{\sin^2 x}$$

$$g'(x) = -\frac{\cos x}{\sin^2 x}$$

$$= -\frac{\cos x}{\sin x} \cdot \frac{1}{\sin x} \Rightarrow g'(x) = -\cot x \csc x$$

$$y' = (2 \sin 2x)(\cos 2x) \cos 2x + \sin^2 2x (-2 \sin 2x)$$

$$y' = 4 \sin 2x \cos^2 2x - 2 \sin^3 2x$$

$$= 2 \sin 2x (2 \cos^2 2x - \sin^2 2x)$$

5) Find the local minimum point on the curve  $y = 2x - \tan x$ , where  $x$  is between negative  $\pi/2$  and  $\pi/2$ .

Turning point, so  $y' = 0$ .

$$y = 2x - \tan x$$

$$y' = 2 - \sec^2 x$$

$$0 = 2 - \sec^2 x$$

$$\sec^2 x = 2$$

$$\frac{1}{\cos^2 x} = 2$$

$$\frac{1}{2} = \cos^2 x$$

on the interval  $\rightarrow \pm \frac{1}{\sqrt{2}} = \cos x$  (related acute  $\rightarrow \frac{\pi}{4}$ )

$$x = \pm \frac{\pi}{4}$$

