Wednesday, October 10, 2018

8.1 Vector and Parametric Equations of a Line in R^2

What forms of an equation of a line do we already know?

$y = mx + b$ → slope y-intercept

$Ax + By + C = 0$ → standard form

Any point on the line will satisfy all of those equations. These forms only work in two dimensions though. In order to extend equations into three space, we need to look at new forms of the equation of a line.

1) Vector Equation of a Line in a Plane

If we have a line with a slope of 3/2 that passes through A(-1, 4), we can also describe it as a line that moves in the same direction as the direction vector $\vec{d} = (2, 3)$. Any scalar multiple of d is also a direction vector.



In general, the vector equation of a line in a plane is:

$$\vec{r} = (x_o, y_o) + t(a, b)$$

where $\vec{r} = (x, y)$ is the position vector of any point on the line, (x_o, y_o) is the position vector of a particular point on the line, and (a, b) is the direction vector, and $t \in R$.

Example: Determine the vector equation of a line with a slope of 3 that passes through the point (2, -5).

$\vec{r} = (2, -5) + t(1, 3)$

Point satisfies both equations.

Note: If you are asked to identify if lines are coincident, you are just checking to see if their direction vectors are parallel.

2) Parametric Equations of a Line

Consider the vector equation $(x, y) = (x_o, y_o) + t(a, b)$. From this, we can see that $x = x_o + ta$ and $y = y_o + tb$. These are called the parametric equations of the line.

Example: Write the equation of the line from the previous example in parametric form.

$\vec{r} = (2, -5) + t(1, 3)$

$x = 2 + t$ $\quad y = -5 + 3t$

The value of t is called a parameter. This is just an independent variable whose value may be chosen. The values of x and y depend on the parameter that is chosen. What does changing the value of t impact with respect to the direction vector? Changing its magnitude

Example: Find the vector and parametric equations of a line that passes through (4, 6) that is perpendicular to $\vec{r} = (2, 1) + t(8, -4)$.

$(x, y) = (4, 6) + t(1, 2)$

$x = 4 + t$ $\quad y = 6 + 2t$

$m = -\frac{4}{8}$ or $\frac{1}{2}$

$m_\perp = \frac{2}{1}$

Practice Problems:

1. Write the vector and parametric equations for the line containing P(-1, 5) and Q(6, 11). Where does this line cross the x - axis?

$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 11}{-1 - 6} = \frac{-6}{-7} = \frac{6}{7}$

$\vec{d} = (7, 6)$

$(x, y) = (-1, 5) + t(7, 6)$

$x = -1 + 7t$

$x = -1 + 7\left(-\frac{5}{6}\right)$

$= -1 - \frac{35}{6}$

$= -\frac{41}{6}$

x-int, so let $y = 0$.

$y = 5 + 6t$

$0 = 5 + 6t$

$-\frac{5}{6} = t$ Use this to find the x-intercept.

$\left(-\frac{41}{6}, 0\right)$

2. A line is defined by the parametric equations $x = -3 - t$ and $y = 5 + 4t$. Does R(4, 21) lie on this line?

$x = -3 - t$ $\quad y = 5 + 4t$

$4 = -3 - t$ $\quad 21 = 5 + 4t$

$7 = -t$ $\quad 16 = 4t$

$-7 = t$ $\quad 4 = t$

These are not equal, so (4, 21) is not on the line.